

## Sturm Liouville Eigenvalue Problems

### Sturm Liouville Differential Operator

$$Lu \equiv -(p(x)u')' + q(x)u \quad x \in (a, b)$$

### Boundary Conditions (homogeneous)

$$B_1 u \equiv \alpha_1 u(a) + \beta_1 u'(a) = 0$$

$$B_2 u \equiv \alpha_2 u(b) + \beta_2 u'(b) = 0$$

Boundary conditions are said to be homogeneous if  $B_k u = 0$  otherwise they are nonhomogeneous as in  $B_k u = 1$ .

Dirichlet Boundary condition has  $\beta_k = 0$  as in

$$B_1 u = u(a) = 0$$

is a Dirichlet B.C. at  $x=a$ .

Neumann Boundary condition has  $\alpha_k = 0$  as in

$$B_2 u = u'(b) = 0$$

If a Boundary condition is neither Dirichlet or Neumann it is said to be a Mixed B.C.

## Sturm Liouville Problem (SLP) Defn

$$(1) \quad Lu = -(p(x)u')' + q(x)u = \lambda u$$

$$(2) \quad B_1 u = \alpha_1 u(a) + \beta_1 u'(a) = 0$$

$$(3) \quad B_2 u = \alpha_2 u(b) + \beta_2 u'(b) = 0$$

Regardless of the value of  $\lambda$  one soln is

$$u(x) \equiv 0 \quad \forall x \in [a, b]$$

The problem is regular only if all hold:

(i)  $[a, b]$  bounded interval

(ii)  $p, p', q \in C[a, b]$

(iii)  $p \neq 0$  on  $[a, b]$

If it is not regular it is a singular (SLP)

Defn Any value  $\lambda_k$  for which (1)-(3)  
has a non trivial soln  $y_k(x) \neq 0$   
is said to be an eigenvalue  
 $y_k(x)$  is the associated eigenfunction

## Nomenclature Examples

$$-(xu')' + x^2 u = \lambda u$$

$$u(0) = 0$$

$$u(1) = 0$$

$$p = x, q = x^2$$

Dirichlet at  $x=0$

Dirichlet at  $x=1$

$$u'' + u = \lambda u$$

$$u'(0) = 0$$

$$u(1) + 2u'(1) = 0$$

$$p = 1, q = 1$$

Neumann at  $x=0$

Mixed at  $x=1$

## Generalizations for SL operators.

$$Lu = x^2 u'' + 2xu' = -(x^2 u')'$$

is a Sturm-Liouville operator where  $p(x) = x^2$

$$Lu = u'' + u' \neq -(pu'' + p'u')$$

for any  $p(x)$  hence is not an SL operator.  
 why? Would have  $p(x)=1$  but  $p'(x)=0$   
 doesn't match the coefficient of  $u'(x)$ .

Any operator eqn

$$Ly = a_1 y'' + a_2 y' + (a_3 + \lambda) y = 0$$

under the transformations/defns

$$p = \exp\left(\int \frac{a_2}{a_1} dx\right) \quad q = \frac{a_3}{a_1} p \quad r = \frac{p}{a_1}$$

becomes

$$(py')' + (q - \lambda r)y = 0$$

which is identical to our theory when  $r=1$ .

## Operator formulation.

If one defines a domain for the differential operator  $L$  as:

$$D = \{ u \in C^2[a, b] : B_1(u) = B_2(u) = 0 \}$$

then  $L$  maps from  $D$  into  $C[a, b]$

$$L : D \rightarrow C[a, b]$$

Here  $u \in C^2[a, b]$  in  $D$  so  $u''$  makes sense in the defn of

$$Lu = -(pu')' + qu$$

With this construct established the Sturm-Liouville Eigenvalue problem is simply

$$\boxed{Lu = \lambda u \quad u \in D}$$

which is much like a matrix e-val problem

$$Ax = \lambda x \quad x \in \mathbb{R}^n$$

## Theorem (Complete $L^2$ basis - SLP)

The regular (SLP)

$$Lu = \lambda u \quad u \in D$$

has infinitely many eigenvalues  $\lambda_k$  with associated eigenfs  $u_k(x)$  have the following properties

(a)  $\lambda_k \in \mathbb{R} \quad k=1, 2, \dots$

(b)  $\lim_{k \rightarrow \infty} |\lambda_k| = \infty$

(c)  $\lambda_j \neq \lambda_k \Rightarrow \langle u_k, u_j \rangle = 0$

(d) for every  $f(x) \in L^2[a, b]$  there are  $c_k$  such that

$$f(x) = \sum_{n=1}^{\infty} c_n u_n(x)$$

in the  $L^2$  sense.

Basically the eigenvalues are real  
become unbounded and e-fns of  
distinct e-values are orthogonal } plain  
english version

Property (d) is called a "completeness"  
property.

Proof (a)  $\lambda_k \in \mathbb{R}$  in text

Proof (b) Efn's with distinct evals are  $\perp$

Let  $\lambda_1 \neq \lambda_2$  be distinct eigenvalues

$$(1) \quad Ly_1 = \lambda_1 y_1 \quad x \in (a, b)$$

$$(2) \quad Ly_2 = \lambda_2 y_2$$

hence

$$(3) \quad \int_a^b (y_2 Ly_1 - y_1 Ly_2) dx = (\lambda_1 - \lambda_2) \int_a^b y_1 y_2 dx$$

$\langle y_1, y_2 \rangle$

Given  $Ly = -(py')' + qy$  eqn (3) becomes

$$(4) \quad (\lambda_1 - \lambda_2) \langle y_1, y_2 \rangle = \int_a^b -y_2 (py_1')' + y_1 (py_2')' dx$$

exact differential

Using the identity

$$\frac{d}{dx} [P(y_1 y_2' - y_2 y_1')] = -y_2 (py_1')' + y_1 (py_2')'$$

Thus (4) becomes

$$(\lambda_1 - \lambda_2) \langle y_1, y_2 \rangle = P(y_1 y_2' - y_2 y_1') \Big|_{x=a}^{x=b}$$

If  $y_k(a) = y_k(b) = 0$  or  $y'_k(a) = y'_k(b) = 0$  the RHS clearly vanishes. Harder to show for Mixed B.C. but

$$(\lambda_1 - \lambda_2) \langle y_1, y_2 \rangle = 0$$

EXAMPLE

Find all eigenvalues and eigenfs  
of the following (SLP)

$$(1) \quad y'' + \lambda y = 0 \quad y(0) = 0 \quad y(\pi) = 0$$

Depending on if  $\lambda < 0$ ,  $\lambda = 0$ ,  $\lambda > 0$  we have  
three different general solutions

$$(2) \quad y(x) = c_1 \sinh \mu x + c_2 \cosh \mu x \quad \lambda = -\mu^2 < 0$$

$$(3) \quad y(x) = c_1 x + c_2 \quad \lambda = 0$$

$$(4) \quad y(x) = c_1 \sin \mu x + c_2 \cos \mu x \quad \lambda = \mu^2 > 0$$

We show that for  $\lambda \leq 0$  only trivial solns  
exist ( $c_1 = c_2 = 0$ ) hence no eigenvalues.

For  $\lambda > 0$  we show trivial solutions unless  
 $\lambda = \lambda_n = n^2$  (eigenvalues).

CASE  $\lambda < 0$  Using general soln (2) above

$$y(0) = c_2 \cosh \mu = 0$$

$$y(\pi) = c_1 \sinh \mu \pi + c_2 \cosh \mu \pi = 0$$

Since both  $\sinh z$  and  $\cosh z$  are positive  
the only solution of the above eqns are

$$c_1 = 0 \quad c_2 = 0$$

hence the (SLP) has no negative e-values.

CASE  $\lambda = 0$  Use general soln (3)

$$y(0) = c_2 = 0$$

$$y(\pi) = c_1\pi + c_2 = 0$$

whose only soln is  $c_1 = c_2 = 0$ . Thus  $\lambda = 0$  is not an e-value of the (SLP)

CASE  $\lambda > 0$  Use general soln (4)

$$y(0) = c_2 = 0$$

$$y(\pi) = c_1 \sin \mu\pi + c_2 \cos \mu\pi = 0$$

Since  $c_2 = 0$  the latter reduces to

$$\sin \mu\pi = 0 \quad \mu_n = n$$

which is true if  $\mu = 1, 2, 3, \dots$  i.e.,  $\mu = \mu_n = n$

Eigenvalues and functions for  $\lambda > 0$

$$\lambda = \lambda_n = \mu_n^2 = n^2$$

with associated eigenfunctions

$$y_n(x) = \sin(nx)$$

$$\lambda_n = n^2$$

Necessarily  $\{y_n\}$  is an orthogonal set on  $[0, \pi]$ .

EXAMPLE Purely Neumann B.C.

$$(1) \quad y'' + \lambda y = 0 \quad y'(0) = y'(\pi) = 0$$

CASE  $\lambda = 0$  General solution  $y(x) = c_1 x + c_2$   
 Since  $y'(x) = c_1$ , must have  $c_1 = 0$

$$\lambda_0 = 0 \quad y_0(x) \equiv 1$$

where the function  $y_0(x) \equiv 1$  is an efn for  $\lambda_0 = 0$ .

CASE  $\lambda = \mu^2 > 0$  General solution

$$y(x) = c_1 \sin(\mu x) + c_2 \cos(\mu x)$$

$$y'(x) = c_1 \mu \cos(\mu x) - c_2 \mu \sin(\mu x)$$

Boundary conditions can be written in matrix form

$$\begin{aligned} y'(0) = 0 & \quad \begin{bmatrix} \mu & 0 \\ \mu \cos(0) & -\mu \sin(0) \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \\ y'(\pi) = 0 & \quad \underbrace{\begin{bmatrix} \mu & 0 \\ \mu \cos(\pi\mu) & -\mu \sin(\pi\mu) \end{bmatrix}}_{A \in \mathbb{R}^{2 \times 2}} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \end{aligned}$$

The system  $A \vec{c} = \vec{0}$  has nontrivial solutions only if

$$\det A = -\mu^2 \sin(\pi\mu) = 0 \quad \left. \begin{array}{l} \text{eval} \\ \text{eqn} \end{array} \right\}$$

from which we deduce  $\mu = \mu_n = n$  (integer).

Also, since  $y'(0) = \mu c_1 = 0 \Rightarrow c_1 = 0$  the eval/e fn pairs are

$$\lambda_n = \mu_n^2 = n^2 \quad y_n(x) = \cos(nx)$$

Collectively

$$\lambda_0 = 0 \quad y_0(x) \equiv 1$$

$$\lambda_n = n^2 \quad y_n(x) = \cos(nx)$$

form a (complete) orthogonal basis for  $L^2[0, \pi]$  in the sense that  $\forall f \in L^2[a, b]$  there exist  $c_k$  such that

$$f(x) = c_0 y_0(x) + \sum_{n=1}^{\infty} c_n y_n(x)$$

in the  $L^2$ -sense.

Detail  $\lambda = -\mu^2 < 0$  not possible since

$$y(x) = c_1 \cosh \mu x + c_2 \sinh \mu x$$

leads to  $A\vec{c} = \vec{0}$  where  $\vec{c} = (c_1, c_2)^T$  and

$$A = \begin{bmatrix} 0 & \mu \\ \mu \sinh \mu \pi & \mu \cosh \mu \pi \end{bmatrix}$$

and

$$\det A = -\mu^2 \sinh \mu \pi \neq 0 \quad \forall \mu.$$

so no negative eigenvalues.

EXAMPLE Eigenvalues of

$$(1) \quad y'' + \lambda y = 0$$

$$(2) \quad y(0) - y'(0) = 0 \quad y(\pi) - y'(\pi) = 0$$

Unlike previous problems  $\lambda < 0$  is possible

$$\text{CASE } \lambda = -\mu^2 < 0 \quad \mu \neq 0$$

$$y(x) = c_1 \cosh \mu x + c_2 \sinh \mu x \quad \text{gen soln}$$

$$y'(x) = c_1 \mu \sinh \mu x + c_2 \mu \cosh \mu x$$

If we use these to evaluate B.C.

$$BC_1 = c_1 + \mu c_2 = 0$$

$$BC_2 = (\cosh \mu \pi - \mu \sinh \mu \pi) c_1 + (\mu \cosh \mu \pi - \sinh \mu \pi) c_2 = 0$$

Written in matrix form this system is

$$A \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} 1 & \mu \\ \cosh \mu \pi - \mu \sinh \mu \pi & \mu \cosh \mu \pi - \sinh \mu \pi \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = 0$$

Non trivial solutions exist only if  $\det A = 0$   
Here

$$\det A = (\mu^2 - 1) \sinh \mu = 0$$

has a soln for  $\mu = \pm 1$ . In either case.  
 $\lambda = -\mu^2 = -1 < 0$  is a negative eval.

For  $\mu = -1$ ,  $BC_1 = 0 \Rightarrow c_1 = c_2$  so that

$$y(x) = c_1 (\cosh \mu x + \sinh \mu x) \quad \mu = -1$$

Using properties of  $\cosh x, \sinh x$  this simplifies to

$$\lambda = -1 \quad y(x) = c_1 e^x$$

as an eval/efn pair.

### CASE $\lambda = 0$

$$y(x) = c_1 x + c_2 \quad \text{gen soln}$$

$$y'(x) = c_1$$

In a similar way the BC yield a system of eqns for  $c_1$  and  $c_2$

$$BC_1: \quad -c_1 + c_2 = 0$$

$$BC_2: \quad (\pi - 1)c_1 + c_2 = 0$$

In matrix form is  $A\vec{c} = 0$  where  $\vec{c} = (c_1, c_2)^T$

$$A = \begin{bmatrix} -1 & 1 \\ (\pi - 1) & 1 \end{bmatrix}$$

Since  $\det A = -\pi \neq 0$  all solns are trivial,  
i.e. have  $c_1 = c_2 = 0$

CASE  $\lambda > 0$   $\lambda = +\mu^2 \quad \mu \neq 0$

$$y(x) = c_1 \cos \mu x + c_2 \sin \mu x$$

$$y'(x) = -c_1 \mu \sin \mu x + c_2 \mu \cos \mu x$$

In a similar fashion the BC yield  
a system of eqns for  $c_k$

$$A \vec{c} = \vec{0} \quad \vec{c} = \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}$$

where (after calculations) we find

$$A = \begin{bmatrix} 1 & -\mu \\ \cos(\mu\pi) + \mu \sin(\mu\pi) & (\sin \mu\pi) - \mu \cos \mu\pi \end{bmatrix}$$

Nontrivial solns are possible only if

$$\det A = (1+\mu^2) \sin(\mu\pi) = 0$$

This eigenvalue eqn  $\Rightarrow \mu_n = n, n = 1, 2, 3$ .  
And from A

$$c_1 - \mu_n c_2 = 0$$

so that  $c_1 = \mu_n c_2$  hence

$$\lambda_n = n^2$$

$$y_n(x) = (n \cos nx + \sin nx)$$

Expansions of  $f(x) \in L^2[0, \pi]$  using these orthogonal functions have the form

$$f(x) = c_0 e^x + \sum_{n=1}^{\infty} c_n \{ n \cos nx + \sin nx \}$$

Summary of evals / efns

$$\lambda_0 = -1$$

$$\phi_0(x) = e^x$$

$$\lambda_n = n^2$$

$$\phi_n(x) = n \cos nx + \sin nx$$

where  $\phi_0(x), \phi_1(x), \phi_2(x), \dots$  are orthogonal under the inner product

$$\langle f, g \rangle = \int_0^\pi f(x)g(x) dx.$$