

Math 450 (2009) – Midterm 1 (Take home)

Due: October 23, 2009.

NAME: _____

1. [30pts] A fluid of density ρ exerts a (drag) force F on a cylinder of diameter D as it flows around it. The fluid has viscosity μ and the fluid velocity far from the cylinder is v . Assume the physical law

$$f(\rho, F, D, \mu, v) = 0$$

and then find all the dimensionless π of the form

$$\pi = \rho^{\alpha_1} F^{\alpha_2} D^{\alpha_3} \mu^{\alpha_4} v^{\alpha_5}$$

Note that the units of viscosity are $[\mu] = ML^{-1}T^{-1}$.

2. [20pts] In quantum mechanics the wave function $\psi(X)$ of a particle of mass m is a solution of Schrödinger's equation. For the quantum harmonic oscillator problem, the (time-independent) Schrödinger equation is:

$$-\frac{\hbar^2}{2m} \frac{d^2\psi}{dX^2} + \frac{1}{2}m\omega^2 X^2\psi = E\psi$$

where X is position, ω is the frequency of the oscillator (potential well), \hbar is Planck's constant and E is the energy of the system. The wave function ψ depends on position X but since it is a probability density function (used to determine the location of the particle) it has no units. Note here that $[\omega] = T^{-1}$ where T is time.

i) Determine the units of \hbar in terms of M, T, L

ii) Nondimensionalize the problem so it has only one dimensionless quantity, i.e.

$$-\frac{d^2\psi}{dx^2} + (x^2 - \mathcal{E})\psi = 0$$

3. [25pts] The equation

$$f(x, \epsilon) = x - \frac{1}{(x + \epsilon)^3} = 0 \quad , \quad 0 < \epsilon \ll 1$$

has two real roots $\bar{x}_{\pm}(\epsilon)$. Assume

$$\bar{x}(\epsilon) = x_0 + x_1\epsilon + O(\epsilon^2)$$

and then determine x_0 and x_1 for both roots. If you use the Binomial theorem note

$$\begin{aligned} (x + \epsilon)^p &= (x_0 + \epsilon x_1 + \dots + \epsilon)^p \\ &= (x_0 + \epsilon(x_1 + 1) + O(\epsilon^2))^p \\ &= x_0^p \left(1 + \epsilon \frac{(x_1 + 1)}{x_0} + O(\epsilon^2) \right)^p \end{aligned}$$

Then you can expand since the latter term has the form $(1 + z)^p$.

4. [25pts] Consider the perturbed first order initial value problem:

$$\frac{dy}{dx} + y^2 = \epsilon \left(\frac{1}{y} - x \right) \quad , \quad y(0) = 1$$

where $0 < \epsilon \ll 1$. Find $y_0(x)$ and $y_1(x)$ in the assumed expansion of the solution y :

$$y(x, \epsilon) = y_0(x) + \epsilon y_1(x) + O(\epsilon^2)$$

Here the leading $O(1)$ problem for $y_0(x)$ is a separable (and Bernoulli).